

1-21-2008

# Long monochromatic Berge cycles in colored 4-uniform hypergraphs

András Gyárfás

*Computer and Automation Research Institute, Hungarian Academy of Sciences, gyarf@szta.hu*

Gábor N. Sárközy

*Worcester Polytechnic Institute, gsarkozy@cs.wpi.edu*

Endre Szemerédi

*Rutgers University - New Brunswick/Piscataway, szemer@cs.rutgers.edu*

Follow this and additional works at: <https://digitalcommons.wpi.edu/computerscience-pubs>



Part of the [Computer Sciences Commons](#)

---

## Suggested Citation

Gyárfás, András, Sárközy, Gábor N., Szemerédi, Endre (2008). Long monochromatic Berge cycles in colored 4-uniform hypergraphs.

.

Retrieved from: <https://digitalcommons.wpi.edu/computerscience-pubs/34>

# Long monochromatic Berge cycles in colored 4-uniform hypergraphs

András Gyárfás\*

Computer and Automation Research Institute  
Hungarian Academy of Sciences  
Budapest, P.O. Box 63  
Budapest, Hungary, H-1518  
[gyarfas@sztaki.hu](mailto:gyarfas@sztaki.hu)

Gábor N. Sárközy†

Computer Science Department  
Worcester Polytechnic Institute  
Worcester, MA, USA 01609  
[gsarkozy@cs.wpi.edu](mailto:gsarkozy@cs.wpi.edu)  
and

Computer and Automation Research Institute  
Hungarian Academy of Sciences  
Budapest, P.O. Box 63  
Budapest, Hungary, H-1518

Endre Szemerédi

Computer Science Department  
Rutgers University  
New Brunswick, NJ, USA 08903  
[szemered@cs.rutgers.edu](mailto:szemered@cs.rutgers.edu)

January 21, 2008

---

\*Research supported in part by OTKA Grant No. K68322.

†Research supported in part by the National Science Foundation under Grant No. DMS-0456401 and by OTKA Grant No. K68322.

### Abstract

Here we prove that for  $n \geq 140$ , in every 3-coloring of the edges of  $K_n^{(4)}$  there is a monochromatic Berge cycle of length at least  $n - 10$ . This result sharpens an asymptotic result obtained earlier. Another result is that for  $n \geq 15$ , in every 2-coloring of the edges of  $K_n^{(4)}$  there is a 3-tight Berge cycle of length at least  $n - 10$ .

## 1 Introduction

Let  $\mathcal{H}$  be an  $r$ -uniform hypergraph (a family of some  $r$ -element subsets of a set). The *shadow graph* of  $\mathcal{H}$  is defined as the graph  $\Gamma(\mathcal{H})$  on the same vertex set, where two vertices are adjacent if they are covered by at least one edge of  $\mathcal{H}$ . A coloring of the edges of an  $r$ -uniform hypergraph  $\mathcal{H}$ ,  $r \geq 2$ , induces a multicoloring on the edges of the shadow graph  $\Gamma(\mathcal{H})$  in a natural way; every edge  $e$  of  $\Gamma(\mathcal{H})$  receives the color of all hyperedges containing  $e$ . We shall denote by  $c(x, y)$  the color set of the edge  $xy$  in  $\Gamma(\mathcal{H})$ . A subgraph of  $\Gamma(\mathcal{H})$  is *monochromatic* if the color sets of its edges have a nonempty intersection. Let  $K_n^{(r)}$  denote the complete  $r$ -uniform hypergraph on  $n$  vertices.

In any  $r$ -uniform hypergraph  $\mathcal{H}$  for  $2 \leq t \leq r$  we define an  $r$ -uniform  $t$ -tight Berge-cycle of length  $\ell$ , denoted by  $C_\ell^{(r,t)}$ , as a sequence of distinct vertices  $v_1, v_2, \dots, v_\ell$ , such that for each set  $(v_i, v_{i+1}, \dots, v_{i+t-1})$  of  $t$  consecutive vertices on the cycle, there is an edge  $e_i$  of  $\mathcal{H}$  that contains these  $t$  vertices and the edges  $e_i$  are all distinct for  $i, 1 \leq i \leq \ell$  where  $\ell + j \equiv j$ . This notion was introduced in [5] and for  $t = 2$  we get ordinary Berge-cycles ([1]) and for  $t = r$  we get the tight cycle (see e.g. [11] or [15]). A Berge-cycle of length  $n$  in a hypergraph of  $n$  vertices is called a Hamiltonian Berge-cycle. It is important to keep in mind that, in contrast to the case  $r = t = 2$ , for  $r > t \geq 2$  a Berge-cycle  $C_\ell^{(r,t)}$ , is not determined uniquely, it is considered as an arbitrary choice from many possible cycles with the same triple of parameters.

In this paper, continuing investigations from [5], [6], [8] and [9], we study long Berge-cycles in hypergraphs. In [5] (by generalizing an earlier conjecture from [6]) the following conjecture was formulated.

**Conjecture 1.1.** *For any fixed  $2 \leq c, t \leq r$  satisfying  $c + t \leq r + 1$  and sufficiently large  $n$ , if we color the edges of  $K_n^{(r)}$  with  $c$  colors, then there is a monochromatic Hamiltonian  $t$ -tight Berge-cycle.*

In [5] it was proved that if the conjecture is true it is best possible, since for any values of  $2 \leq c, t \leq r$  satisfying  $c + t > r + 1$  the statement is not true. The conjecture was proved for  $r = 3$  in [6]. The asymptotic form of the conjecture was proved for  $r = 4$  and  $t = 2$  in [6] and for every  $r$  and  $t = 2$  in [9] - in both papers the Regularity Lemma was used. In this paper we apply an elementary approach and we study the  $r = 4$  case. We prove the conjecture in both cases ( $c = 3, t = 2$  and  $c = 2, t = 3$ ) with a constant error term.

**Theorem 1.2.** *Suppose that an 3-coloring is given on the edges of  $K_n^{(4)}$ , where  $n \geq 140$ . Then there is a monochromatic Berge-cycle of length at least  $n - 10$ .*

This sharpens the asymptotic result obtained earlier for  $r = 4$  in [6].

**Theorem 1.3.** *Suppose that an 2-coloring is given on the edges of  $K_n^{(4)}$ , where  $n \geq 15$ . Then there is a monochromatic 3-tight Berge-cycle of length at least  $n - 10$ .*

## 2 Proofs

**Proof of Theorem 1.2.** Suppose that  $c$  is a 3-coloring on the edges of  $\mathcal{K} = K_n^{(4)}$ , where  $n \geq 140$ . Color  $i \in c(x, y)$  on the edge  $xy$  of  $G = \Gamma(\mathcal{K})$  is a *good color* if at least 3 edges of color  $i$  contain  $\{x, y\}$  in  $\mathcal{K}$ . We consider  $G$  with a new coloring  $c^*$  where  $c^*(x, y) \subseteq c(x, y)$  is the set of good colors on  $xy$ . Assuming that  $\binom{n-2}{2} > 6$ , i.e.  $n > 6$ , every edge of  $\mathcal{K}$  has at least one color in  $c^*$ .

Suppose first that some edge  $xy$  of  $G = \Gamma(\mathcal{K})$  is colored (under  $c^*$ ) with a single color, say with color 1. We claim that there is a Hamiltonian Berge cycle in  $\mathcal{K}$  in color 1. Indeed, the definition of  $xy$  implies that under  $c^*$  at most four edges of  $H = G \setminus \{x, y\}$  are not colored with 1. Since for  $n > 10$  we have  $n - 6 > (n - 2)/2$ , the color 1 subgraph of  $H$  satisfies Dirac's condition (see [13]), and thus one can easily find a Hamiltonian path  $P = \{y_1, \dots, y_{n-2}\}$  of color 1 in  $H$  such that there are two extra edges  $y_1 y_p$  and  $y_{n-2} y_k$  of color 1 from the endpoints of  $P$  with  $2 < p, k < n - 3$ . Now the cyclic ordering  $x, y_1, y_2, \dots, y_{n-2}, y$  defines a Hamiltonian Berge-cycle in color 1 with the following edge assignments. For  $x, y_1$  assign  $e_n = \{x, y_1, y_p, y\}$ . For  $y_j, y_{j+1}$  ( $1 \leq j \leq n - 3$ ) assign  $e_j = \{x, y, y_j, y_{j+1}\}$ , for  $y_{n-2}, y$  assign  $e_{n-2} = \{y_{n-2}, y, y_k, x\}$ , and finally for  $x, y$  we can assign  $e_{n-1}$  as any edge of color 1 containing  $x, y$  and different from all other  $e_i$ -s.

Now we may assume that  $c^*$  colors all edges of  $G$  with one of the four color sets: 12, 13, 23, 123.

**Lemma 2.1.** *Assume that there is a monochromatic Hamiltonian cycle  $C$  in  $G$  under coloring  $c^*$ . Then there is a Hamiltonian Berge-cycle in  $\mathcal{K}$  under coloring  $c$ .*

*Proof.* Assume that  $C = x_1, x_2, \dots, x_n$  is a Hamiltonian cycle of  $G$  in color 1 (under  $c^*$ ). Then, following the cyclic order of vertices on  $C$ , let  $A_j$  be the set of edges of  $\mathcal{K}$  in color 1 containing  $x_j, x_{j+1}$ . Since each  $A_j$  has at least three elements and no element of  $A_j$  covers more than three consecutive pairs of  $C$ , Hall's theorem ensures a one-to one correspondence from the consecutive pairs to the sets  $A_j$ . This clearly defines the required Hamiltonian Berge-cycle.  $\square$

We need some observations on the structure of the coloring  $c^*$ . Let  $x$  be an arbitrary vertex, define  $U_{12}(x), U_{13}(x), U_{23}(x), U_{123}(x)$  as the sets to which  $x$  is connected in color sets 12, 13, 23, 123 respectively. Define

$$B_i = \{x \in V(G) \mid U_{ij} = U_{ik} = \emptyset, U_{jk} \neq \emptyset\},$$

where  $i, j, k$  are the elements of  $\{1, 2, 3\}$  in some order. Observe that the  $B_i$ -s are pairwise disjoint, within the  $B_i$ -s every edge of  $G$  has color set  $\{j, k\}$

or 123, and for  $j \neq i$ , an edge of  $G$  from  $B_i$  to  $B_j$  has color set 123. Set  $B_4 = \{x \in V(G) \mid |U_{123}(x)| \geq n/2\}$ .

**Lemma 2.2.** *Suppose that  $\cup_{i=1}^4 B_i = V(G)$ . Then there is a Hamiltonian cycle  $G$  in the coloring  $c^*$ .*

*Proof.* Suppose w.l.o.g that  $|B_1| \leq |B_2| \leq |B_3|$ . We show that there is a Hamiltonian cycle in color 1. Denoting the degree of a vertex  $v$  in color  $i$  by  $d_i(v)$ , we have that  $d_1(v) \geq |B_2| + |B_3| \geq |B_2| + |B_1|$  if  $v \in B_1$ ,  $d_1(v) = n-1$  if  $v \in B_2 \cup B_3$  and  $d_1(v) \geq \frac{n}{2}$  if  $v \notin \cup_{i=1}^3 B_i$  (since in the latter case  $v \in B_4$ ). These conditions immediately imply - through either Pósa's or Chvátal's condition (see [13]) that there is a Hamiltonian cycle.  $\square$

Thus we may assume that there exists  $x \in V(G) \setminus \cup_{i=1}^4 B_i$  (otherwise Lemmas 2.1 and 2.2 would finish the proof). Set  $U = V(G) \setminus (\{x\} \cup U_{123})$  and assume w.l.o.g.  $|U_{23}| \leq |U_{12}| \leq |U_{13}|$ . Since  $x \notin B_2$  we have  $U_{12} \neq \emptyset$  and  $x \notin B_4$  implies that  $|U| \geq \lfloor n/2 \rfloor$ .

We show that  $|U_{23}| \leq 1$ . Indeed, otherwise we may select two two-element sets  $A_{23} \subseteq U_{23}, A_{12} \subseteq U_{12}$  and a five-element set  $A_{13} \subseteq U_{13}$ . (The condition  $|U| \geq \lfloor n/2 \rfloor$  implies that  $|U_{13}| \geq \frac{\lfloor n/2 \rfloor}{3} \geq 5$  so  $A_{13}$  can be defined.) For every fixed  $u_{23} \in A_{23}$  there are at most two edges of color 1 among the edges of  $\mathcal{K}$  in the form  $\{x, u_{23}, x_{12}, x_{13}\}$  where  $x_{12} \in A_{12}, x_{13} \in A_{13}$  are arbitrary. Repeating this argument for fixed  $u_{12}, u_{13}$  we get that there are at most  $4 + 4 + 10 = 18$  edges of  $\mathcal{K}$  in the form  $\{x, u_{23}, x_{12}, x_{13}\}$ . However, there are  $2 \times 2 \times 5 = 20$  such edges giving a contradiction.

Now we fix  $y \in U_{12}, z \in U_{13}$  and define a graph  $H$  on the vertices of  $V(G) \setminus (U_{23} \cup \{x, y, z\})$  as follows. Let  $uv \in E(H)$  be an edge of  $H$  in the following cases: (i)  $u \in U_{13}, c(\{x, y, u, v\}) = 1$ , in this case the edge is called an  $xy$ -edge; (ii)  $u \in U_{12}, c(\{x, z, u, v\}) = 1$ , now the edge is called an  $xz$ -edge. Set  $|V(H)| = N$  and note that  $N \geq n - 4$ .

**Lemma 2.3.** *The graph  $H$  has a cycle  $C$  of length at least  $N - 6$  in color 1.*

*Proof.* Set

$$T_{12} = U_{12} \cap V(H), T_{13} = U_{13} \cap V(H), T = U \cap V(H), T_{123} = U_{123}.$$

Consider an arbitrary vertex  $u \in T_{12} \cup T_{13}$ . Set  $w = z$  if  $u \in T_{12}$  otherwise set  $w = y$ . Apart from at most four choices of  $v \in V(H)$  the edge  $\{x, u, w, v\}$  of  $\mathcal{K}$  is of color 1. Thus every vertex of  $T \subseteq V(H)$  has degree at least  $N - 5$  in  $H$ . Consider the set  $S \subseteq T_{123}$  of vertices whose degrees are at most 11 in the bipartite subgraph  $[T, T_{123}]$  of  $H$ . Observe that

$$|T|(|T_{123}| - 5) \leq |E[T, T_{123}]| \leq (|T_{123}| - |S|)|T| + 11|S|$$

implying that  $|S| \leq 6$  if  $66 \leq |T|$  and this is true since  $|T| > \lfloor n/2 \rfloor - 4 > 65$ . Now consider the subgraph  $F$  of  $H$  induced by  $T \cup (T_{123} \setminus S)$ . In fact, we may assume that  $|S| = 6$  since deleting  $6 - |S|$  vertices does not influence the following observation: each vertex  $v \in T$  has degree at least  $N - 11$  in  $F$  and each vertex

$v \in T_{123} \setminus S$  has degree more than 11. Now we can apply Chvátal's condition (see [13]) to prove that there is a Hamiltonian cycle in  $F \subset H$ . Indeed, with  $M = |V(F)|$ , we have to show that  $d_k \leq k < \frac{M}{2}$  implies that  $d_{M-k} \geq M - k$  where  $d_1 \leq d_2 \leq \dots \leq d_M$  is the degree sequence of  $F$ . This is immediate because the number of vertices with possibly small degrees (i.e.  $v \in T_{123} \setminus S$ ) is at most

$$|U_{123}| - 6 \leq \left\lfloor \frac{n}{2} \right\rfloor - 6 \leq \left\lfloor \frac{N+4}{2} \right\rfloor - 6 = \left\lfloor \frac{M+10}{2} \right\rfloor - 6 = \left\lfloor \frac{M}{2} \right\rfloor - 1. \quad (1)$$

Indeed, let us take a  $k$  for which  $d_k \leq k < \frac{M}{2}$ .  $11 < d_k \leq k$  implies that  $k > 11$ . But then from (1) we get

$$d_{M-k} \geq d_{\lceil \frac{M}{2} \rceil} \geq N - 11 \geq M - 11 > M - k,$$

as desired.  $\square$

To finish the proof of Theorem 1.2, observe that the cycle  $C$  obtained from Lemma 2.3 defines a Berge-cycle if its  $xy$ -edges and  $xz$ -edges are extended (with  $\{x, y\}$  or with  $\{x, z\}$  to edges of  $\mathcal{K}$ . Thus we have a Berge-cycle of length  $N - 6 \geq n - 10$  as required.  $\square$

**Proof of Theorem 1.3.** Suppose that a 2-coloring  $c$  is given on the edges of  $\mathcal{K} = K_n^{(4)}$ . Let  $V$  be the vertex set of  $\mathcal{K}$  and observe that  $c$  defines a 2-multicoloring on the complete 3-uniform hypergraph  $\mathcal{T}$  with vertex set  $V$  by coloring a triple  $T$  with the colors of the edges of  $\mathcal{K}$  containing  $T$ . We say that  $T \in \mathcal{T}$  is *good in color  $i$*  if  $T$  is contained in at least two edges of  $\mathcal{K}$  of color  $i$  ( $i = 1, 2$ ).

**Lemma 2.4.** *Every edge  $xy \in E(G)$  is in at least  $n - 4$  good triples of the same color.*

*Proof.* Consider an edge  $xy$  in  $G$ . Coloring  $c$  induces a 2-coloring  $c'$  on  $W = V \setminus \{x, y\}$ . Applying a result of Bollobás and Gyárfás, [2], there exists a subgraph  $H$  with at least  $|W| - 2 = n - 4$  vertices such that  $H$  is 2-connected and monochromatic under  $c'$ , say in color 1. In particular, every vertex of  $H$  has degree at least two in color 1. Thus, for every vertex  $z$  of  $H$ ,  $\{x, y, z\}$  is a good triple in color 1.  $\square$

Using Lemma 2.4, we can define a 2-coloring  $c^*$  on the shadow graph  $G = \Gamma(\mathcal{K})$  by coloring  $xy \in E(G)$  with the color of the (at least  $n - 4$ ) good triples containing  $xy$ . Using a well-known result about the Ramsey number of even cycles ([4], [14]) there is a monochromatic even cycle  $C$  of length  $2t$  where  $2t = \lceil \frac{2n}{3} \rceil - 6$  or  $2t = \lceil \frac{2n}{3} \rceil - 7$ . (In fact there is a bit longer cycle, but that is too long for our purposes.) Assume that  $C$  is in color 1. Label the edges of  $C$  as  $e_j = \{p_j, p_{j+1}\}$ ,  $j = 1, 2, \dots, 2t$ . We use here index arithmetic  $\pmod{2t}$ .

We shall find a large Berge-cycle in color 1 with the following greedy procedure. By Lemma 2.4, for each  $i \in [2t]$  there is a set  $A_i \subset V$  such that  $|A_i| \geq n - 4$  and the triple  $T_i = \{p_i, p_{i+1}, x\}$  is good in color 1 for every  $x \in A_i$ . We claim

that we can find a set  $\{v_j \in A_{2j-1} \setminus V(C)\}$  for  $j \in [t]$  with the following property: for every  $j \in [t]$ ,

$$v_j \in A_{2j-2} \cap A_{2j-1} \cap A_{2j}.$$

Assume that for  $j \leq h < t$  we have this property and there are at least seven vertices in  $S = V \setminus (V(C) \cup \{\cup_{j=1}^h v_j\})$ . Indeed, if  $|S| \geq 7$ , then - because each of the three sets intersects  $S$  in at least five elements -  $U = S \cap A_{2h} \cap A_{2h+1} \cap A_{2h+2} \neq \emptyset$  so we can select  $v_{h+1} \in U$ . Now we only have to observe that during the whole process

$$|S| \geq n - 3t \geq n - \frac{3}{2}(\lceil \frac{2n}{3} \rceil - 6) \geq 7,$$

and thus the claim is proved.

Now we finish the proof by claiming that the cyclic permutation  $P = p_1, v_1, p_2, p_3, v_2, p_4, \dots, p_{2t-1}, v_t, p_1$  determines a Berge-cycle. Indeed, from the definition of  $v_j$ , every triple of three consecutive vertices on  $P$  is good in color 1. Therefore at least two edges  $\mathcal{K}$  of color 1 are available to cover a consecutive triple. However, no edge of  $\mathcal{K}$  can cover more than two consecutive triples of  $P$ . Thus, by Hall's theorem, there is a matching from the consecutive triples of  $P$  to the set of color 1 edges of  $\mathcal{K}$  containing them. The length of this Berge-cycle is  $3t \geq \frac{3}{2}(\lceil \frac{2n}{3} \rceil - 7) \geq n - 10$ .  $\square$

## References

- [1] C. Berge, *Graphs and Hypergraphs*, North Holland, Amsterdam and London, 1973.
- [2] B. Bollobás, A. Gyárfás, Highly connected monochromatic subgraphs, to appear in *Discrete Mathematics*.
- [3] A. Figaj, T. Łuczak, The Ramsey number for a triple of long even cycles, *Journal of Combinatorial Theory, Ser. B*, **97** (2007), pp. 584-596.
- [4] R.J. Faudree, R.H. Schelp, All Ramsey numbers for cycles in graphs, *Discrete Mathematics* **8** (1974), pp. 313-329.
- [5] P. Dorbec, S. Gravier, G.N. Sárközy, Monochromatic Hamiltonian  $t$ -tight Berge-cycles in hypergraphs, submitted for publication.
- [6] A. Gyárfás, J. Lehel, G.N. Sárközy, R. H. Schelp, Monochromatic Hamiltonian Berge-cycles in colored complete uniform hypergraphs, accepted for publication in the *Journal of Combinatorial Theory, Ser. B*.
- [7] A. Gyárfás, M. Ruszinkó, G.N. Sárközy, E. Szemerédi, Three-color Ramsey numbers for paths, *Combinatorica*, **27** (2007), pp. 35-69.
- [8] A. Gyárfás, G.N. Sárközy, The 3-color Ramsey number of a 3-uniform Berge-cycle, submitted for publication.

- [9] A. Gyárfás, G.N. Sárközy, E. Szemerédi, Monochromatic matchings in the shadow graph of almost complete hypergraphs, submitted for publication.
- [10] P. Haxell, T. Łuczak, Y. Peng, V. Rödl, A. Ruciński, M. Simonovits, J. Skokan, The Ramsey number for hypergraph cycles I, *Journal of Combinatorial Theory, Ser. A* 113 (2006), pp. 67-83.
- [11] P. Haxell, T. Łuczak, Y. Peng, V. Rödl, A. Ruciński, J. Skokan, The Ramsey number for hypergraph cycles II, manuscript.
- [12] T. Łuczak,  $R(C_n, C_n, C_n) \leq (4 + o(1))n$ , *Journal of Combinatorial Theory, Ser. B* 75 (1999), pp. 174-187.
- [13] L. Lovász, *Combinatorial Problems and Exercises*, 2. edition, North-Holland, 1979.
- [14] V. Rosta, On a Ramsey type problem of Bondy and Erdős, I and II, *Journal of Combinatorial Theory B* 15 (1973), pp. 94-120.
- [15] V. Rödl, A. Ruciński, E. Szemerédi, A Dirac-type theorem for 3-uniform hypergraphs, *Combinatorics, Probability and Computing* 15 (2006), pp. 229-251.