

April 2018

# Analysis of Polyelectrolytes and Functionalized Silica for Precious Metal Catalyst Recovery

Amy Elizabeth Toscano  
*Worcester Polytechnic Institute*

Patrick M. Bresnahan  
*Worcester Polytechnic Institute*

Follow this and additional works at: <https://digitalcommons.wpi.edu/mqp-all>

---

## Repository Citation

Toscano, A. E., & Bresnahan, P. M. (2018). *Analysis of Polyelectrolytes and Functionalized Silica for Precious Metal Catalyst Recovery*. Retrieved from <https://digitalcommons.wpi.edu/mqp-all/1851>

This Unrestricted is brought to you for free and open access by the Major Qualifying Projects at Digital WPI. It has been accepted for inclusion in Major Qualifying Projects (All Years) by an authorized administrator of Digital WPI. For more information, please contact [digitalwpi@wpi.edu](mailto:digitalwpi@wpi.edu).

# Analysis of Polyelectrolytes and Functionalized Silica for Precious Metal Catalyst Recovery

A Major Qualifying Project

Submitted to the Faculty

of

WORCESTER POLYTECHNIC INSTITUTE

in partial fulfillment of the requirements for the

Degree of Bachelor of Science

by

---

Patrick Bresnahan

---

Amy Toscano

Date: 4/26/2018

Sponsoring Organization: Sunovion Pharmaceuticals

Project Advisors:

---

Professor Stephen Kmiotek, Advisor

---

Professor Drew Brodeur, Co-Advisor

*This report represents work of WPI undergraduate students submitted to the faculty as evidence of a degree requirement. WPI routinely publishes these reports on its web site without editorial or peer review. For more information about the projects program at WPI, see*

<http://www.wpi.edu/Academics/Projects>.

## **Abstract**

*Sunovion Pharmaceuticals utilizes an organometallic rhodium-based catalyst in one of their drug production processes. Due to its high cost, it is beneficial to recover for reuse; however, due to the low concentration of the catalyst in the waste stream of the process, it has proven difficult to separate from solution. Our team investigated the use of polyelectrolytes and functionalized silica metal scavengers as a means to recover the catalyst. Through the use of the ICP-OES, rhodium concentrations were analyzed before and after the addition of the polyelectrolytes and metal scavengers. While the polyelectrolytes proved ineffective due to the lack of precipitation and inability to filter, the functionalized silica metal scavengers showed over 90% recovery of rhodium at various loadings.*

This MQP contains information deemed confidential to the business interest of the industrial sponsor. Please contact Stephen Kmiolek at [sjkmiolek@wpi.edu](mailto:sjkmiolek@wpi.edu) or Drew Brodeur at [dbrodeur@wpi.edu](mailto:dbrodeur@wpi.edu) for additional information.